

Asymptotic symmetries on Kerr–Newman horizon without anomaly of diffeomorphism invariance

Jun-ichirou Koga

Advanced Research Institute for Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555, Japan

E-mail: koga@gravity.phys.waseda.ac.jp

Abstract. We analyze asymptotic symmetries on the Killing horizon of the four-dimensional Kerr–Newman black hole. We first derive the asymptotic Killing vectors on the Killing horizon, which describe the asymptotic symmetries, and find that the general form of these asymptotic Killing vectors is the universal one possessed by arbitrary Killing horizons. We then construct the phase space associated with the asymptotic symmetries. It is shown that the phase space of an extreme black hole either has the size comparable with a non-extreme black hole, or is small enough to exclude degeneracy, depending on whether or not the global structure of a Killing horizon particular to an extreme black hole is respected. We also show that the central charge in the Poisson brackets algebra of these asymptotic symmetries vanishes, which implies that there is not an anomaly of diffeomorphism invariance. By taking into account other results in the literature, we argue that the vanishing central charge on a black hole horizon, in an effective theory, looks consistent with the thermal feature of a black hole. We furthermore argue that the vanishing central charge implies that there are infinitely many classical configurations that are associated with the same macroscopic state, while these configurations are distinguished physically.

1. Introduction

In spite of various efforts, the statistical origin of the thermal feature of a black hole [1, 2] still remains to be clarified. A universal framework will be necessary to understand, from a general point of view, what are microscopic degrees of freedom responsible for the thermal feature and how those degrees of freedom constitute a thermal object. One of possible ideas is to consider a universal and local geometric structure associated with existence of a black hole horizon and analyze whether such a geometric structure is related to the thermal feature of a black hole. Motivated by the success in the case of the B.T.Z. black hole [3], asymptotic symmetries on a black hole horizon have been analyzed in this context (see [4, 5] for recent reviews), but it still remains controversial whether this approach is successful, in particular whether the desirable form of a non-vanishing central charge, i.e., an anomaly of diffeomorphism invariance, arises *naturally* in the algebra associated with the asymptotic symmetries, as in the case of the B.T.Z. black hole.

An idea behind the asymptotic symmetries on a black hole horizon is the possibility that the microscopic degrees of freedom responsible for the thermal feature may be described without the details of quantum theory of gravity. This idea does not seem absurd, because the microscopic degrees of freedom of *standard* thermal radiation are indeed described within the classical electromagnetism. Whereas the quantization condition should be imposed in order to obtain the Planckian spectrum, we need not resort to the complete theory of quantum physics to study the thermodynamical state of standard thermal radiation. It is then expected that we may understand also the microscopic degrees of freedom responsible for the thermal feature of a black hole in the context of classical theory. In the case of a black hole, however, we are faced with the fact that the structure of a Killing horizon, which is fundamental in black hole thermodynamics, does not allow for a sufficient number of classical configurations, while a large number of microscopic states are necessary for a black hole to exhibit the thermal feature and possess the entropy. Then, a possible way is to consider a weakened structure of a Killing horizon, and analyze whether it allows for a sufficient number of classical configurations.

In the previous work [6], a local geometric structure of a Killing horizon, called an asymptotic Killing horizon, was analyzed from the viewpoint of universality, by weakening the geometric structure of a Killing horizon without assuming any global structures of a spacetime or field equations. An asymptotic Killing horizon was defined as the pair (H, ξ^a) of a null hypersurface H and its generator ξ^a , i.e., the vector field that plays the same role only on H as the generator of a Killing horizon. The generator of an asymptotic Killing horizon was then found to be given by the asymptotic Killing vectors, which describe the asymptotic symmetries on an asymptotic Killing horizon. It was thus shown that once there exists one asymptotic Killing horizon, there exist infinitely many asymptotic Killing horizons (H, ξ^a) on the same null hypersurface H , that is, the generator ξ^a is highly non-unique. These results show that we can indeed obtain degeneracy of classical configurations by weakening the structure of a Killing horizon. We also showed that asymptotic Killing horizons are physically distinguishable, not being a sort of gauge, by analyzing the behavior of the acceleration associated with the generators ξ^a . We furthermore argued that the discrepancy between string theory [7, 8, 9] and the Euclidean approach to black hole thermodynamics [10, 11] in the entropy of an extreme black hole will be resolved, if the microscopic states responsible for the thermal feature of a black hole are connected with the asymptotic Killing horizons.

It then may be expected that the microscopic states of black hole thermodynamics are described by asymptotic Killing horizons. To clarify whether this is true or not, however, it is necessary to understand how asymptotic Killing horizons are described in a phase space, especially when we are interested in quantum and/or statistical physics of asymptotic Killing horizons. In particular, we need to derive the Poisson brackets algebra of the asymptotic symmetries in order to understand whether a non-vanishing central charge arises. It is important also to show that asymptotic Killing horizons can be regarded as degenerate from a macroscopic point of view, and hence as the microscopic states that constitute one thermodynamical state. An evidence for this will be provided, if we can show that the same values of thermodynamical variables are shared by all the asymptotic Killing horizons. In order to analyze these issues, however, we should specify the explicit form of the Lagrangian of the theory we consider. We will thus focus in this paper on the four-dimensional Einstein–Maxwell theory, and hence on the Kerr–Newman black hole spacetime, as an example that is basic but interesting enough. By considering explicitly the metric near the Killing horizon of the Kerr–Newman black hole, we can derive also the general form of the asymptotic Killing vectors on the Killing horizon. Thus, the purpose of this paper is to investigate those aspects associated with the asymptotic Killing vectors which are clarified by specifying the black hole solution and the Lagrangian, and provide a further evidence that the microscopic states of black hole thermodynamics are described by asymptotic Killing horizons.

To analyze the asymptotic Killing vectors, it is reasonable to focus on regular and continuous asymptotic symmetries, unless convincing physical explanations are provided, which show that singularity or discontinuity of asymptotic symmetries is essential in black hole thermodynamics. Then, we will utilize in this paper a regular coordinate system to derive the asymptotic Killing vectors in a correct and well-defined manner, while results never depend on a coordinate system used in the analysis. In the next section, we will thus first review the behavior of the metric and the gauge potential of the electromagnetic field, in a regular coordinate system and a regular gauge, near the Killing horizon of the Kerr–Newman black hole. In section 3, we will derive the asymptotic Killing vectors on the Killing horizon, based on the behavior of the metric considered in section 2. It will be found that the general form of the asymptotic Killing vectors so derived is the universal one possessed by arbitrary Killing horizons [6]. We then construct in section 4 the phase space associated with the asymptotic symmetries and compute the Poisson brackets between the conserved charges conjugate to the asymptotic Killing vectors. To do so, we will need to incorporate the electromagnetic field into the asymptotic symmetries, and it will help to observe an interesting feature of the phase space of an extreme black hole. From the computation of the Poisson brackets, we will see that the central charge in the Poisson brackets algebra vanishes, and that this fact itself implies that all asymptotic

Killing horizons give the same value of a thermodynamical quantity. Finally, we will summarize and discuss our results in section 5.

2. Near horizon

The metric $\bar{g}_{ab}$ and the gauge potential $\bar{A}_a^s$ of the four-dimensional Kerr–Newman black hole are written in the Boyer–Lindquist coordinate system (t, r, θ, φ) as

$$d\bar{s}^2 = -\frac{\Delta - a^2 \sin^2 \theta}{\Sigma} dt^2 - 2 \frac{a \sin^2 \theta (r^2 + a^2 - \Delta)}{\Sigma} dt d\varphi + \frac{(r^2 + a^2)^2 - \Delta a^2 \sin^2 \theta}{\Sigma} \sin^2 \theta d\varphi^2 + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2, \quad (2.1)$$

$$\bar{A}_a^s = -\frac{Qr}{\Sigma} (dt)_a + \frac{aQr \sin^2 \theta}{\Sigma} (d\varphi)_a, \quad (2.2)$$

where the bar on a field variable indicates that it is used in the following sections as a background, and the superscript s on $\bar{A}_a$ implies that it is singular. The functions Δ and Σ are defined by

$$\Delta \equiv r^2 - 2Mr + a^2 + Q^2 \equiv (r - r_+)(r - r_-), \quad \Sigma \equiv r^2 + a^2 \cos^2 \theta, \quad (2.3)$$

and M , $J \equiv aM$, and Q are the mass, the angular momentum, and the electric charge of the black hole, while the constants r_+ and r_- are defined as

$$r_{\pm} \equiv M \pm \sqrt{M^2 - a^2 - Q^2}, \quad (2.4)$$

and denote the radius of the outer and the inner horizon, respectively. We focus in this paper on the case of $r_+ \geq r_-$, so that the spacetime singularity is hidden behind an event horizon.

In the Kerr–Newman spacetime $(M, \bar{g}_{ab})$, there exist the timelike Killing vector $\xi_{(t)}^a \equiv (\partial/\partial t)^a$ and the rotational Killing vectors $\xi_{(i)}^a$. If the black hole is not rotating ($a = 0$), $\xi_{(i)}^a$ denotes all the Killing vectors that generate $O(3)$ rotations, whereas only the axial Killing vector $\xi_{(\varphi)}^a \equiv (\partial/\partial \varphi)^a$ is allowed in the rotating case ($a \neq 0$). The linear combination of these Killing vectors defined by

$$\xi_{(h)}^a \equiv \xi_{(t)}^a + \Omega_H \xi_{(\varphi)}^a, \quad (2.5)$$

where $\Omega_H \equiv a/(r_+^2 + a^2)$ is the angular velocity of the black hole, becomes null on $r = r_+$, and so it generates a Killing horizon on $r = r_+$. The field strength $\bar{F}_{ab} = \bar{\nabla}_a \bar{A}_b - \bar{\nabla}_b \bar{A}_a$ of the electromagnetic field also is invariant along these Killing vectors as

$$\mathcal{L}_{\xi_{(p)}} \bar{F}_{ab} = 0, \quad (2.6)$$

where $\xi_{(p)}^a$ denotes collectively the Killing vectors $\xi_{(h)}^a$ and $\xi_{(i)}^a$. Although the Lie derivatives of the gauge potential along $\xi_{(p)}^a$ need not vanish in general, the gauge potential $\bar{A}_a^s$ given by (2.2) satisfies $\mathcal{L}_{\xi_{(p)}} \bar{A}_a^s = 0$ for all the Killing vectors $\xi_{(p)}^a$.

Focusing on the future horizon, it is convenient to consider here a double null coordinate system that is “co-rotating” with the future horizon. (One can of course consider the past horizon in the same manner.) We thus define the new coordinates U , V and ϕ by[‡]

$$dt = dV - \frac{r^2 + a^2}{\Delta} dr, \quad \frac{r^2 + a^2}{\Delta} dr = \frac{1}{2}(dV - dU), \quad d\varphi = d\phi + \Omega_H dV - \frac{a}{\Delta} dr. \quad (2.7)$$

However, the coordinate system (U, V, θ, ϕ) does not cover the Killing horizon. In the case of a non-extreme black hole $r_+ > r_-$, we then define the Kruskal coordinates u and v as

$$u = -\frac{1}{\kappa_0} \exp[-\kappa_0 U], \quad v = \frac{1}{\kappa_0} \exp[\kappa_0 V], \quad (2.8)$$

[‡] The coordinate system defined by (2.7) differs from the standard one (see, e.g., [17]) in the second term on the right-hand side of the last equation, which is given here by $\Omega_H dV$, not $\Omega_H dt$. However, it is convenient to use the coordinate system adopted here, when we consider the metric near the future horizon of an extreme black hole.

where $\kappa_0 \equiv (r_+ - r_-)/2(r_+^2 + a^2)$ is the surface gravity of the Killing horizon. Now, the future horizon is located at $u = 0$ and covered by this coordinate system smoothly. Indeed, near the future horizon $u = 0$, the metric $\bar{g}_{ab}$ behaves as

$$d\bar{s}^2 = -2\Lambda du dv + 2\Xi v du d\phi + \Sigma_+ d\theta^2 + \frac{R^4}{\Sigma_+} \sin^2 \theta d\phi^2 + O(u), \quad (2.9)$$

where the functions Λ , Ξ , and Σ_+ are defined by

$$\Lambda \equiv \frac{1}{2} \exp[-2\kappa_0 r_+] \left(1 - \frac{r_-}{r_+}\right)^{\alpha+1} \frac{r_+^2}{r_+^2 + a^2} (1 - a\Omega_H \sin^2 \theta), \quad (2.10)$$

$$\Xi \equiv \frac{a\kappa_0}{2} \exp[-2\kappa_0 r_+] \left(1 - \frac{r_-}{r_+}\right)^{\alpha+1} \frac{r_+^2}{r_+^2 + a^2} \sin^2 \theta, \quad \Sigma_+ \equiv r_+^2 + a^2 \cos^2 \theta, \quad (2.11)$$

and the constants R and α are given by

$$R^2 \equiv r_+^2 + a^2, \quad \alpha \equiv \frac{r_-^2 + a^2}{r_+^2 + a^2}. \quad (2.12)$$

As one can see from the components in the regular coordinate system (u, v, θ, ϕ) , the gauge potential $\bar{A}_a^S$ is singular on the future horizon $u = 0$. A regular gauge is then achieved by the gauge transformation $\bar{A}_a^S \rightarrow \bar{A}_a^N = \bar{A}_a^S - \bar{\nabla}_a \bar{\chi}^N$ with

$$\bar{\chi}^N \equiv -\frac{Qr_+}{(r_+^2 + a^2)} t = -\frac{Qr_+}{2(r_+^2 + a^2)} V - \frac{Qr_+}{2(r_+^2 + a^2)} U, \quad (2.13)$$

and the new gauge potential $\bar{A}_a^N$ is written near $u = 0$ as

$$\bar{A}_a^N = O(u^0) (du)_a + \frac{Qr}{\Sigma} a \sin^2 \theta (d\phi)_a + O(u). \quad (2.14)$$

For all the Killing vectors $\xi_{(p)}^a$, we can show

$$\mathcal{L}_{\xi_{(p)}} \bar{A}_a^N = -\bar{\nabla}_a \mathcal{L}_{\xi_{(p)}} \bar{\chi}^N = 0, \quad (2.15)$$

and hence the gauge potential $\bar{A}_a^N$ also obeys the isometries of the Kerr–Newman black hole. Moreover, we have

$$\xi_{(h)}^c \bar{A}_c^N = O(u), \quad (2.16)$$

where $\xi_{(h)}^a$ is written in the present coordinate system as

$$\xi_{(h)}^a = \kappa_0 v \left(\frac{\partial}{\partial v} \right)^a - \kappa_0 u \left(\frac{\partial}{\partial u} \right)^a. \quad (2.17)$$

We can obtain (2.16) from (2.14) and (2.17). However, it is important to note that (2.16) follows essentially because $\bar{A}_a^N$ is regular and $\xi_{(h)}^a$ vanishes on the bifurcation surface ($u = v = 0$) of the Killing horizon [18], and (2.15) yields $\mathcal{L}_{\xi_{(h)}} \xi_{(h)}^c \bar{A}_c^N = 0$. In other words, (2.16) results from the fact that a non-extreme black hole has a bifurcate Killing horizon.

When the black hole is extreme $r_+ = r_-$, we introduce null coordinates u and v by

$$U = -2r_+ f(-r_+^{-1}u), \quad V = v, \quad (2.18)$$

where the function $f(x)$ is defined as

$$f(x) \equiv x + 2 \ln x - \frac{\gamma^2}{x}, \quad (2.19)$$

and the constant γ is given by

$$\gamma^2 \equiv \frac{R^2}{r_+^2} = 1 + \frac{a^2}{r_+^2}. \quad (2.20)$$

The future horizon is found to be located at $u = 0$ in this coordinate system, and the metric $\bar{g}_{ab}$ behaves smoothly near $u = 0$ as

$$d\bar{s}^2 = -2\Lambda du dv + 2\Xi du d\phi + \Sigma_+ d\theta^2 + \frac{R^4}{\Sigma_+} \sin^2 \theta d\phi^2 + O(u), \quad (2.21)$$

where Λ and Ξ are defined in the extreme case as

$$\Lambda \equiv 1 - a \Omega_H \sin^2 \theta, \quad \Xi \equiv a \sin^2 \theta, \quad (2.22)$$

while Σ_+ and R are defined by (2.11) and (2.12), respectively, as in the non-extreme case. Also in the case of an extreme black hole, the gauge potential $\bar{A}_a^s$ is singular on $u = 0$, and we obtain a regular gauge potential $\bar{A}_a^t$ by performing the gauge transformation $\bar{A}_a^s \rightarrow \bar{A}_a^t = \bar{A}_a^s - \bar{\nabla}_a \bar{\chi}^t$, where $\bar{\chi}^t$ is given as

$$\bar{\chi}^t \equiv -\frac{Q r_+}{2(r_+^2 + a^2)} U - \frac{Q r_+}{2(r_+^2 + a^2)} V - Q r_+^2 \frac{r_+^2 - a^2}{(r_+^2 + a^2)^2} \rho(-r_+^{-1}u) + \frac{Q}{2} \frac{r_+^2 - a^2}{(r_+^2 + a^2)^2} u v, \quad (2.23)$$

and $\rho(x)$ is defined by

$$\rho(x) \equiv \frac{1}{2}x^2 + 2x + \gamma^2 \ln x. \quad (2.24)$$

(One might expect that a regular gauge is achieved by the same gauge transformation (2.13) as in the non-extreme case, since the gauge potential takes the same form when expressed in the coordinate system (U, V, θ, ϕ) . However, it cannot remove all the singular parts of $\bar{A}_a^s$ in the case of an extreme black hole.)

The new gauge potential $\bar{A}_a^t$ behaves near $u = 0$ as

$$\bar{A}_a^t = O(u^0) (du)_a + \frac{Q r_+}{\Sigma_+} a \sin^2 \theta (d\phi)_a + O(u). \quad (2.25)$$

However, the gauge transformation $\bar{A}_a^s \rightarrow \bar{A}_a^u = \bar{A}_a^s - \bar{\nabla}_a \bar{\chi}^u$, where

$$\bar{\chi}^u \equiv -\frac{Q r_+}{2(r_+^2 + a^2)} U - Q r_+^2 \frac{r_+^2 - a^2}{(r_+^2 + a^2)^2} \rho(-r_+^{-1}u) + \frac{Q}{2} \frac{r_+^2 - a^2}{(r_+^2 + a^2)^2} u v, \quad (2.26)$$

also yields a regular gauge potential $\bar{A}_a^u$, which is written near $u = 0$ as

$$\bar{A}_a^u = -\frac{Q r_+}{2(r_+^2 + a^2)} (dv)_a + O(u^0) (du)_a + \frac{Q r_+}{\Sigma_+} a \sin^2 \theta (d\phi)_a + O(u). \quad (2.27)$$

Therefore, both of the gauge potentials, $\bar{A}_a^t$ and $\bar{A}_a^u$, are regular all over the future horizon. We also note that we have

$$\mathcal{L}_{\xi_{(h)}} \bar{A}_a = -\bar{\nabla}_a \mathcal{L}_{\xi_{(h)}} \bar{\chi} = O(u), \quad (2.28)$$

where $\bar{A}_a$ and $\bar{\chi}$ denote $\bar{A}_a^t$ and $\bar{\chi}^t$, or $\bar{A}_a^u$ and $\bar{\chi}^u$, respectively, and $\xi_{(h)}^a$ is expressed in the present coordinate system as

$$\xi_{(h)}^a = \left(\frac{\partial}{\partial v} \right)^a + \frac{1}{2} \frac{u^2}{(u - r_+)^2 + a^2} \left(\frac{\partial}{\partial u} \right)^a. \quad (2.29)$$

Since the Lie derivatives of $\bar{A}_a^t$ and $\bar{A}_a^u$ along the rotational Killing vectors $\xi_{(i)}^a$ vanish, all the isometries in the Kerr–Newman spacetime are respected by both of $\bar{A}_a^t$ and $\bar{A}_a^u$ at least on the future horizon $u = 0$ as

$$\mathcal{L}_{\xi_{(p)}} \bar{A}_a^t = O(u), \quad \mathcal{L}_{\xi_{(p)}} \bar{A}_a^u = O(u). \quad (2.30)$$

On the other hand, the difference between these two gauge potentials, which is important below, is described as

$$\xi_{(h)}^c \bar{A}_c^t = O(u), \quad \xi_{(h)}^c \bar{A}_c^u = -\frac{Q r_+}{2(r_+^2 + a^2)} + O(u). \quad (2.31)$$

We note that $\xi_{(h)}^c \bar{A}_c^t$ vanishes on the future horizon $u = 0$, whereas $\xi_{(h)}^c \bar{A}_c^u$ does not. Thus, $\bar{A}_a^t$ possesses the same property as $\bar{A}_a^N$ in the non-extreme case, which satisfies (2.16), while $\bar{A}_a^u$ does not possess this property. Although $\xi_{(h)}^c \bar{A}_c^N$ in the non-extreme case is required to vanish on the horizon in a regular gauge, the corresponding quantity need not vanish in the case of an extreme black hole. This is because the Killing horizon of an extreme black hole does not have a bifurcation surface. Then, there are no reasons to prefer to the gauge $\bar{A}_a^t$ and exclude the gauge $\bar{A}_a^u$. Rather, it is $\bar{A}_a^u$ that reflects faithfully the global geometric structure of a Killing horizon particular to an extreme black hole, while $\bar{A}_a^t$ imitates the gauge in the non-extreme case.

3. Asymptotic Killing vectors

In the previous work [6], an asymptotic Killing vector on an asymptotic Killing horizon was defined based on the feature of the standard asymptotic symmetries, such as the B.M.S. group [12] and the conformal group in a 3-dimensional asymptotically anti-de Sitter spacetime [13], where the leading behavior of the asymptotic metrics is left unchanged under the asymptotic symmetry transformations, no matter what their sub-leading behavior is. In particular, an asymptotic Killing vector on a Killing horizon H in a spacetime $(M, \bar{g}_{ab})$ is defined to be a smooth vector ζ^a that satisfies

$$\mathcal{L}_\zeta g_{ab} = O(\sigma), \quad (3.1)$$

for any smooth scalar σ such that $\sigma = 0$ and $\bar{\nabla}_a \sigma \neq 0$ on H , where g_{ab} are arbitrary smooth metrics that coincide with $\bar{g}_{ab}$ on H as

$$g_{ab} = \bar{g}_{ab} + O(\sigma). \quad (3.2)$$

Thus, an asymptotic Killing vector behaves as if a Killing vector only on H , and hence it is a generalized notion of a Killing vector. We note also that the infinitesimal diffeomorphisms $g_{ab} \rightarrow g_{ab} + \mathcal{L}_\zeta g_{ab}$ along asymptotic Killing vectors ζ^a leave the form (3.2) of g_{ab} unchanged, which therefore generate the asymptotic symmetry group.

We now derive explicitly the general form of the asymptotic Killing vectors ζ^a on the future Killing horizon of the Kerr–Newman black hole, by solving (3.1). To do so, here we specifically take, as the scalar function σ in (3.1), the retarded null coordinates u introduced by (2.8) in the non-extreme case and (2.18) in the extreme case, because these functions are smooth, vanishing, and non-degenerate on the future horizon. We then substitute the asymptotic form of the metric, (2.9) in the non-extreme case and (2.21) in the extreme case, for $\bar{g}_{ab}$ in (3.2), and expand the asymptotic Killing vector ζ^a as

$$\zeta^a = \zeta_{(0)}^a(v, \theta, \phi) + u \zeta_{(1)}^a(v, \theta, \phi) + O(u^2). \quad (3.3)$$

The requirement that (3.1) holds for arbitrary perturbed metrics g_{ab} of the form (3.2) gives ten equations for the components of ζ^a , and they are classified into two sets of equations, which decouple from each other, and one equation given as

$$\zeta_{(0)}^u = 0. \quad (3.4)$$

Among these two sets of equations, the first set consists of equations for the leading order of the angular components $\zeta_{(0)}^\theta$ and $\zeta_{(0)}^\phi$, whose explicit forms are given by

$$\partial_v \zeta_{(0)}^\theta = 0, \quad (3.5)$$

$$\partial_v \zeta_{(0)}^\phi = 0, \quad (3.6)$$

$$\zeta_{(0)}^\theta \partial_\theta \Sigma_+ + 2\Sigma_+ \partial_\theta \zeta_{(0)}^\theta = 0, \quad (3.7)$$

$$\Sigma_+ \partial_\phi \zeta_{(0)}^\theta + \frac{R^4}{\Sigma_+} \sin^2 \theta \partial_\theta \zeta_{(0)}^\phi = 0, \quad (3.8)$$

$$\zeta_{(0)}^\theta \partial_\theta \left(\frac{R^4}{\Sigma_+} \sin^2 \theta \right) + 2 \frac{R^4}{\Sigma_+} \sin^2 \theta \partial_\phi \zeta_{(0)}^\phi = 0, \quad (3.9)$$

in both the non-extreme and the extreme cases. We note from (3.5) and (3.6) that $\zeta_{(0)}^\theta$ and $\zeta_{(0)}^\phi$ do not depend on v , and hence (3.7)–(3.9) are purely two-dimensional equations. Indeed, they are nothing but the components of the two-dimensional Killing equation on a horizon sphere (a cross-section of the future Killing horizon) with v fixed, and therefore the exact rotational Killing vectors $\xi_{(i)}^a$ satisfy these equations. To see whether or not other solutions (“enhanced symmetries” only at the horizon) happen to exist, we employ the axial symmetry of the Kerr–Newman spacetime and write $\zeta_{(0)}^\theta$ and $\zeta_{(0)}^\phi$ as

$$\zeta_{(0)}^\theta = P(\theta) e^{im\phi}, \quad \zeta_{(0)}^\phi = Q(\theta) e^{im\phi}. \quad (3.10)$$

By substituting (3.10), we rewrite (3.7)–(3.9) as

$$P(\theta) \partial_\theta \Sigma_+ + 2\Sigma_+ \partial_\theta P(\theta) = 0, \quad (3.11)$$

$$i m \Sigma_+^2 P(\theta) + R^4 \sin^2 \theta \partial_\theta Q(\theta) = 0, \quad (3.12)$$

$$P(\theta) \frac{\Sigma_+}{\sin^2 \theta} \partial_\theta \left(\frac{1}{\Sigma_+} \sin^2 \theta \right) = -2 i m Q(\theta), \quad (3.13)$$

respectively. The solution for $m = 0$ is then given by $P(\theta) = 0$ and $Q(\theta) = \text{const.}$, and hence it is expressed, by using $\partial/\partial\varphi = \partial/\partial\phi$, as a constant multiple of the axial Killing vector $\xi_{(\varphi)}^a$. For $m \neq 0$, on the other hand, we integrate (3.11) and substitute it into (3.13), which gives $P(\theta)$ and $Q(\theta)$. When they are substituted further into (3.12), we obtain

$$m^2 \Sigma_+^4(\theta) + (r_+^2 + a^2)^3 \left(3a^2 \sin^2 \theta \cos^2 \theta - \Sigma_+(\theta) \right) = 0, \quad (3.14)$$

unless $P(\theta)$ and $Q(\theta)$ are trivial as $P(\theta) = Q(\theta) = 0$, i.e., $\zeta_{(0)}^\theta = \zeta_{(0)}^\phi = 0$. It is then straightforward to show that non-trivial solutions of $P(\theta)$ and $Q(\theta)$ for $m \neq 0$ do not exist when $a \neq 0$. Actually, (3.14) gives $m^2 = 1$ when evaluated at $\theta = 0$, and it gives $a^2 = 0$ when it is evaluated at $\theta = \pi/2$ and $m^2 = 1$ is substituted. Therefore, a non-trivial solution in the case of $a \neq 0$ is possible only for $m = 0$, and is given by a constant multiple of $\xi_{(\varphi)}^a$. On the other hand, when $a = 0$, the spacetime is spherically symmetric, and hence the three exact Killing vectors of $O(3)$ rotations are allowed. Moreover, in this case, any solutions should be expressed as linear combinations of these three Killing vectors with constant coefficients, because (3.7)–(3.9) are the components of the two-dimensional Killing equation. Thus, in any case, we see that the solutions of $\zeta_{(0)}^\theta$ and $\zeta_{(0)}^\phi$ are described by

$$\zeta_{(0)}^\theta \left(\frac{\partial}{\partial\theta} \right)^a + \zeta_{(0)}^\phi \left(\frac{\partial}{\partial\phi} \right)^a = a^{(i)} \xi_{(i)}^a, \quad (3.15)$$

where the coefficients $a^{(i)}$ are arbitrary constants and summation over the index (i) is understood, if necessary.

On the other hand, the second set of equations is written in the matrix form as

$$\begin{pmatrix} 0 & -\Lambda & 0 & v\Xi \\ -\Lambda & 0 & 0 & 0 \\ 0 & 0 & \Sigma_+ & 0 \\ v\Xi & 0 & 0 & R^4 \sin^2 \theta / \Sigma_+ \end{pmatrix} \begin{pmatrix} \zeta_{(1)}^u \\ \zeta_{(1)}^v \\ \zeta_{(1)}^\theta \\ \zeta_{(1)}^\phi \end{pmatrix} \\ = \begin{pmatrix} 0 \\ \kappa_0 \hat{\omega} \Lambda \\ 0 \\ -\kappa_0 v \hat{\omega} \Xi \end{pmatrix} + \begin{pmatrix} 0 \\ \kappa_0 v \Lambda \partial_v \hat{\omega} \\ \kappa_0 v \Lambda \partial_\theta \hat{\omega} \\ \kappa_0 v \Lambda \partial_\phi \hat{\omega} \end{pmatrix} + \begin{pmatrix} 0 \\ \zeta_{(0)}^\theta \partial_\theta \Lambda \\ -v \Xi \partial_\theta \zeta_{(0)}^\phi \\ -v \zeta_{(0)}^\theta \partial_\theta \Xi - v \Xi \partial_\phi \zeta_{(0)}^\phi \end{pmatrix}, \quad (3.16)$$

in the non-extreme case, and as

$$\begin{pmatrix} 0 & -\Lambda & 0 & \Xi \\ -\Lambda & 0 & 0 & 0 \\ 0 & 0 & \Sigma_+ & 0 \\ \Xi & 0 & 0 & R^4 \sin^2 \theta / \Sigma_+ \end{pmatrix} \begin{pmatrix} \zeta_{(1)}^u \\ \zeta_{(1)}^v \\ \zeta_{(1)}^\theta \\ \zeta_{(1)}^\phi \end{pmatrix} = \begin{pmatrix} 0 \\ \Lambda \partial_v \hat{\omega} \\ \Lambda \partial_\theta \hat{\omega} \\ \Lambda \partial_\phi \hat{\omega} \end{pmatrix} + \begin{pmatrix} 0 \\ \zeta_{(0)}^\theta \partial_\theta \Lambda \\ -\Xi \partial_\theta \zeta_{(0)}^\phi \\ -\zeta_{(0)}^\theta \partial_\theta \Xi - \Xi \partial_\phi \zeta_{(0)}^\phi \end{pmatrix}, \quad (3.17)$$

in the extreme case. The function $\hat{\omega}$ in (3.16) and (3.17) is defined by

$$\hat{\omega}(v, \theta, \phi) \equiv \frac{1}{\kappa_0 v} \zeta_{(0)}^v(v, \theta, \phi), \quad (3.18)$$

in the non-extreme case, and by

$$\hat{\omega}(v, \theta, \phi) \equiv \zeta_{(0)}^v(v, \theta, \phi), \quad (3.19)$$

in the extreme case, while it is left undetermined and hence arbitrary in what follows. We note that it is possible to write (3.18) and (3.19) in a unified manner as

$$\zeta_{(0)}^v = \hat{\omega} \xi_{(h)}^v, \quad (3.20)$$

by using (2.17) and (2.29). We also note from (3.15) that we have $\zeta_{(0)}^\theta = 0$ and $\zeta_{(0)}^\phi = \text{const.}$ in the case of $a \neq 0$. On the other hand, when $a = 0$, we see from (2.10), (2.11) and (2.22) that $\Lambda = \text{const.}$ and

$\Xi = 0$. Thus, in any case, the last terms on the right-hand side of (3.16) and (3.17) vanish. In addition, the components of the matrices on the left-hand side of (3.16) and (3.17) are the leading order of the components of $\bar{g}_{ab}$, and the components of the first term on the right-hand side of (3.16) are given by the leading order of $-\kappa_0 \hat{\omega} \bar{g}_{ab} \delta_u^b$. Then, by multiplying (3.16) and (3.17) by $u \bar{g}^{ca}$, and using the explicit form (2.17) of the Killing vector $\xi_{(h)}^a$ in the non-extreme case, we obtain

$$u \zeta_{(1)}^c = \hat{\omega} \delta_u^c \xi_{(h)}^u + u \kappa_0 v \Lambda \bar{\nabla}^c \hat{\omega} + O(u^2), \quad (3.21)$$

in the non-extreme case, and

$$u \zeta_{(1)}^c = u \Lambda \bar{\nabla}^c \hat{\omega} + O(u^2), \quad (3.22)$$

in the extreme case. Therefore, by collecting (3.4), (3.20), (3.21), and (3.22) all together, adding the independent solution (3.15), and noting from (2.29) that $\xi_{(h)}^u$ in the extreme case is $O(u^2)$, we see that the general solution of ζ^a is written as

$$\zeta^a = \hat{\omega} \xi_{(h)}^a + a^{(i)} \xi_{(i)}^a + u X^a + O(u^2), \quad (3.23)$$

where X^a is given by

$$X^a \equiv \kappa_0 v \Lambda \bar{\nabla}^a \hat{\omega}, \quad (3.24)$$

in the non-extreme case, and by

$$X^a \equiv \Lambda \bar{\nabla}^a \hat{\omega}. \quad (3.25)$$

in the extreme case.

We find that (3.23) coincides with the result in [14] in the spherically symmetric case. It is also possible to rewrite (3.23) into the form presented in [6], which manifestly shows that (3.23) is defined covariantly. To see this, we recall that the normal $\bar{\nabla}_a u$ to the future horizon $u = 0$ does not vanish, and that the future horizon is a null hypersurface. Therefore, $\bar{\nabla}^a u$ is proportional to $\xi_{(h)}^a$ on $u = 0$, and hence there exists a smooth scalar n such that

$$\xi_{(h)}^a = n \bar{\nabla}^a u + O(u) = \bar{\nabla}^a (nu) + O(u). \quad (3.26)$$

Actually, n is given by $n = -\kappa_0 v \Lambda + O(u)$ in the non-extreme case, and $n = -\Lambda + O(u)$ in the extreme case. Then, by defining a scalar λ as $\lambda = nu$, which is found from (3.26) to satisfy $\xi_{(h)}^a = \bar{\nabla}^a \lambda + O(u)$ and hence is called the potential of $\xi_{(h)}^a$ [6], we see from (3.24) and (3.25) that we can write as

$$u X^a = -un \bar{\nabla}^a \hat{\omega} + O(u^2) = -\lambda \bar{\nabla}^a \hat{\omega} + O(u^2), \quad (3.27)$$

in both the non-extreme and the extreme cases. Furthermore, although $\hat{\omega}$ has been introduced in (3.18) and (3.19) as a function of v , θ , and ϕ only, it can indeed depend on u , as well. By expanding an arbitrary smooth function $\omega(u, v, \theta, \phi)$, which depends also on u , into the Taylor series near $u = 0$ as

$$\omega(u, v, \theta, \phi) = \hat{\omega}(v, \theta, \phi) + u \omega'(v, \theta, \phi) + O(u^2), \quad (3.28)$$

and using (3.26), we obtain

$$\omega \xi_{(h)}^a - un \bar{\nabla}^a \omega = \hat{\omega} \xi_{(h)}^a - un \bar{\nabla}^a \hat{\omega} + O(u^2). \quad (3.29)$$

Thus, dependence of ω on u cancels out automatically to linear order in u , and then ζ^a is eventually written as

$$\zeta^a = \omega \xi_{(h)}^a + a^{(i)} \xi_{(i)}^a - \lambda \bar{\nabla}^a \omega + O(u^2). \quad (3.30)$$

This is indeed the form of the asymptotic Killing vectors presented in [6], which was shown to satisfy (3.1) independently of the choice of σ . Therefore, the general form, i.e., the only possible form, of the asymptotic Killing vectors on the Killing horizon of the Kerr–Newman black hole is found to be given by (3.30). An important feature of the asymptotic Killing vectors (3.30) is that they exist universally on arbitrary Killing horizons [6], not only on the Killing horizon of the Kerr–Newman black hole.

4. Covariant phase space

4.1. Symplectic formalism

Although the asymptotic Killing vectors are defined to act on the metric g_{ab} in a spacetime, it is important to consider also how the actions of the asymptotic Killing vectors are represented in a phase space. To analyze this issue, we employ the symplectic (covariant phase space) formalism [19, 20, 21], which we here briefly review.

We write as $L(\psi)$ the Lagrangian of an n -dimensional theory, where ψ^I denotes the dynamical field variables collectively. The actions $\mathcal{L}_\zeta \psi^I$ of diffeomorphisms along arbitrary vectors ζ^a are then identified with the variations $\delta_\zeta \psi^I$ induced by $\mathcal{L}_\zeta \psi^I$ in the covariant phase space, i.e., $\delta_\zeta \psi^I = \mathcal{L}_\zeta \psi^I$. In the covariant phase space, on the other hand, the variations $\delta_\zeta \psi^I$ are generated through the Poisson brackets between ψ^I and the conserved charges $\mathcal{H}[\psi, \zeta]$ conjugate to the vectors ζ^a , which are defined by their derivatives $\delta \mathcal{H}[\psi, \zeta]$ as

$$\delta \mathcal{H}[\psi, \zeta] \equiv \int_{\mathcal{C}} \omega_{c_1 \dots c_{n-1}}(\psi; \delta_\zeta \psi, \delta \psi). \quad (4.1)$$

Here, $\mathcal{C}$ stands for an arbitrary (partial) Cauchy surface, on which the conserved charges $\mathcal{H}[\psi, \zeta]$ are defined, and $\omega_{c_1 \dots c_{n-1}}(\psi; \delta_1 \psi, \delta_2 \psi)$ is the symplectic current density constructed as

$$\omega_{c_1 \dots c_{n-1}}(\psi; \delta_1 \psi, \delta_2 \psi) \equiv \delta_2 \left(\varepsilon_{bc_1 \dots c_{n-1}} \Theta^b(\psi; \delta_1 \psi) \right) - \delta_1 \left(\varepsilon_{bc_1 \dots c_{n-1}} \Theta^b(\psi; \delta_2 \psi) \right), \quad (4.2)$$

from the surface term $\Theta^b(\psi; \delta \psi)$ in the variation

$$\delta \left(\varepsilon_{c_1 \dots c_n} L(\psi) \right) = \varepsilon_{c_1 \dots c_n} E_I \delta \psi^I + \varepsilon_{c_1 \dots c_n} \nabla_b \Theta^b(\psi; \delta \psi), \quad (4.3)$$

of the Lagrangian density $\varepsilon_{c_1 \dots c_n} L(\psi)$, where $E_I = 0$ denotes the field equations.

However, we should impose boundary conditions on ψ^I so that the conserved charges $\mathcal{H}[\psi, \zeta]$ are integrable. The integrability condition of $\mathcal{H}[\psi, \zeta]$ is then written [21] as

$$\int_{\partial \mathcal{C}} \zeta^b \omega_{bc_1 \dots c_{n-2}}(\psi; \delta_1 \psi, \delta_2 \psi) = 0, \quad (4.4)$$

for arbitrary independent variations $\delta_1 \psi^I$ and $\delta_2 \psi^I$ tangent to the covariant phase space. When the integrability condition (4.4) is satisfied, the conserved charges $\mathcal{H}[\psi, \zeta]$ are expressed in the form of a surface integral as

$$\mathcal{H}[\psi; \zeta] = \int_{\partial \mathcal{C}} \frac{1}{2} \varepsilon_{bac_1 \dots c_{n-2}} \left[Q^{ba}(\psi; \zeta) + 2 \zeta^{[b} B^{a]}(\psi) \right] + \mathcal{H}_0[\zeta], \quad (4.5)$$

where $\mathcal{H}_0[\zeta]$ is an integration constant, which we tentatively set to zero, $B^a(\psi)$ is defined by

$$\delta \int_{\partial \mathcal{C}} \varepsilon_{bac_1 \dots c_{n-2}} \zeta^b B^a(\psi) \equiv \int_{\partial \mathcal{C}} \varepsilon_{bac_1 \dots c_{n-2}} \zeta^b \Theta^a(\psi; \delta \psi), \quad (4.6)$$

and $Q^{ba}(\psi; \zeta)$ is given, by using $E_I = 0$, as the potential of the Noether current $J^b(\psi; \zeta) \equiv \Theta^b(\psi; \mathcal{L}_\zeta \psi) - \zeta^b L$, i.e., $J^b(\psi; \zeta) = \nabla_a Q^{ba}(\psi, \zeta)$.

Under the integrability condition (4.4), it has been also shown [14] that the Poisson brackets between $\mathcal{H}[\psi, \zeta]$ are given by

$$\left\{ \mathcal{H}[\psi; \zeta_1], \mathcal{H}[\psi; \zeta_2] \right\} = \mathcal{H}[\psi; \mathcal{L}_{\zeta_1} \zeta_2] + K[\zeta_1, \zeta_2], \quad (4.7)$$

where the central term $K[\zeta_1, \zeta_2]$ is evaluated as

$$K[\zeta_1, \zeta_2] = \delta_{\zeta_2} \mathcal{H}[\bar{\psi}; \zeta_1] - \mathcal{H}[\bar{\psi}; \mathcal{L}_{\zeta_1} \zeta_2], \quad (4.8)$$

on a background configuration $\bar{\psi}^I$, and $\delta_{\zeta_2} \mathcal{H}[\bar{\psi}; \zeta_1]$ is calculated by

$$\begin{aligned} \delta_{\zeta_2} \mathcal{H}[\bar{\psi}; \zeta_1] = \int_{\partial \mathcal{C}} \frac{1}{2} \bar{\varepsilon}_{bac_1 \dots c_{n-2}} \left[\mathcal{L}_{\zeta_2} Q^{ba}(\bar{\psi}; \zeta_1) - \mathcal{L}_{\zeta_1} Q^{ba}(\bar{\psi}; \zeta_2) - Q^{ba}(\bar{\psi}; \mathcal{L}_{\zeta_2} \zeta_1) \right. \\ \left. + Q^{ba}(\bar{\psi}; \zeta_1) (\bar{\nabla}_d \zeta_2^d) - Q^{ba}(\bar{\psi}; \zeta_2) (\bar{\nabla}_d \zeta_1^d) + 2 \zeta_1^{[b} \zeta_2^{a]} L(\bar{\psi}) \right]. \end{aligned} \quad (4.9)$$

The same expression for the central term has been derived by Silva [15]. Although Barnich and Brandt [16] gave a slightly different expression, the difference is shown to vanish, when $\mathcal{L}_\zeta \bar{g}_{ab}$ is vanishing on $\partial\mathcal{C}$, as we consider in this paper.

In order to analyze the integrability condition (4.4) and the central term (4.8) for the actions of the asymptotic Killing vectors on the Killing horizon of the four-dimensional Kerr–Newman black hole, we now let ζ^a be given by (3.30) and $\bar{\psi}^I$ be the Kerr–Newman black hole solution. Correspondingly, we consider the four-dimensional Einstein–Maxwell theory, and thus the Lagrangian $L(\psi)$ is given by

$$L(\psi) = \frac{1}{16\pi} \left[R - F^{ab} F_{ab} \right]. \quad (4.10)$$

As the dynamical field variables ψ^I , we take the metric g_{ab} and the gauge potential A_a , because the variation of the Lagrangian (4.10) with respect to these variables yields the field equations $E_I = 0$, from which we also find that $\Theta^b(\psi; \delta\psi)$ is given as

$$\Theta^b(\psi; \delta\psi) = \frac{1}{16\pi} \left[g_{ac} \nabla^b \delta g^{ac} - \nabla_a \delta g^{ba} - 4F^{ba} \delta A_a \right]. \quad (4.11)$$

The background configuration $\bar{\psi}^I$ is thus described by the metric $\bar{g}_{ab}$, along with the regular gauge potential $\bar{A}_a^N$ in the non-extreme case, and $\bar{A}_a^I$ or $\bar{A}_a^H$ in the extreme case, as we presented in section 2. While we focus in this paper on the spacetime exterior to the horizon of the Kerr–Newman black hole, the boundary $\partial\mathcal{C}$ of $\mathcal{C}$ is considered to consist of a horizon sphere only, by assuming that all the asymptotic Killing vectors fall off rapidly on the other boundary of $\mathcal{C}$ at infinity. In order to complete the construction of the covariant phase space, we now specify the behavior of the metric and the gauge potential near $\partial\mathcal{C}$ so that the integrability condition (4.4) is satisfied.

4.2. Boundary condition

Since the symplectic current density $\omega_{bcd}(\psi; \delta_1\psi, \delta_2\psi)$ is bi-linear in $\delta\psi^I$ or their derivatives, the integrability condition (4.4) is satisfied, if we impose the boundary condition that $\delta\psi^I$ and their derivatives fall off rapidly enough near the horizon sphere $\partial\mathcal{C}$. However, we are analyzing here the actions of the asymptotic symmetries in the covariant phase space, and thus we wish to impose a boundary condition so that any configurations of ψ^I achieved by the asymptotic symmetry transformations reside within the covariant phase space.

The desired boundary condition on the metric g_{ab} is easily specified. We note that (3.1) gives

$$\delta_\zeta g_{ab} = \mathcal{L}_\zeta g_{ab} = \mathcal{O}(u), \quad (4.12)$$

as long as g_{ab} takes the form of (3.2). On the other hand, (3.2) is described in the variational form as

$$\delta g_{ab} = \mathcal{O}(u). \quad (4.13)$$

We note that (4.12) is consistent with (4.13), and hence the actions of all the asymptotic Killing vectors ζ^a are realized within a phase space, when the phase space is constructed under the boundary condition (4.13). In addition, it has been shown [14] that if (4.13) is imposed, the integrability condition (4.4) of the conserved charges is satisfied in vacuum Einstein gravity. Also in presence of the electromagnetic field, we then naturally impose (4.13) as the boundary condition on the metric g_{ab} .

In order to specify the boundary condition on the gauge potential A_a , we first need to incorporate A_a into the notion of the asymptotic symmetries, so that the asymptotic Killing vectors ζ^a act on A_a as the generators of the asymptotic symmetries. Since the background gauge potential $\bar{A}_a$ is invariant along the Killing vectors $\xi_{(p)}^a$ on the horizon, as it is seen from (2.15) and (2.30), and the asymptotic Killing vectors are local generalization of Killing vectors, it might seem reasonable to require that $\bar{A}_a$ is invariant also along the asymptotic Killing vectors ζ^a as $\mathcal{L}_\zeta \bar{A}_a = \mathcal{O}(u)$, and hence $\delta_\zeta \bar{A}_a = \mathcal{O}(u)$. By noting $\zeta^a \nabla_a u = \mathcal{O}(u)$ and following the same argument as we used above for the boundary condition on the metric, it then would be natural to impose the boundary condition on the gauge potential as $\delta A_a = \mathcal{O}(u)$. This would be plausible

also from the viewpoint of integrability of the conserved charges $\mathcal{H}[\psi, \zeta]$, since the integrability condition (4.4) is satisfied when $\delta A_a = O(u)$, as we will see from (4.14) below.

However, the electromagnetic field has the gauge degree of freedom. We thus write as $\delta A_a = \nabla_a \chi + O(u)$, where χ may be arbitrary at this stage but will be constrained from the integrability condition in what follows. By using (4.11) and substituting (3.30) and (4.13) into (4.4), we find that the integrability condition is described as

$$\int_{\partial\mathcal{C}} \varepsilon_{cd} \omega \left[(\delta_1 F^{ba}) \xi_{(h)b} (\delta_2 A_a) - (\delta_2 F^{ba}) \xi_{(h)b} (\delta_1 A_a) \right] = 0, \quad (4.14)$$

where ε_{cd} is the volume element on $\partial\mathcal{C}$ and we note that $\xi_{(i)}^a$ are tangent to $\partial\mathcal{C}$. Since ω is an arbitrary function, it then follows that the inside of the bracket in (4.14) should vanish in order that the integrability condition is satisfied for all the asymptotic Killing vectors. We see that it does vanish if we impose the condition $\chi = O(u)$ near the horizon, because F^{ab} is antisymmetric and $\delta A_a = \nabla_a \chi + O(u) \propto \nabla_a u + O(u) \propto \xi_{(h)a} + O(u)$ in that case. Imposing a boundary condition on χ might look strange, but it is actually natural that a boundary condition on the metric is combined with a condition on gauge transformations on the boundary, as it occurs in the case of an asymptotically anti-de Sitter spacetime [23], for example. Therefore, we impose the boundary condition on A_a as

$$\delta A_a = \nabla_a (u \chi') + O(u), \quad (4.15)$$

and hence A_a is written as

$$A_a = \bar{A}_a + \nabla_a (u \chi') + O(u), \quad (4.16)$$

where χ' is an arbitrary smooth function.

One can see that the boundary condition (4.15) is supported by the natural requirement that the conserved charges $\mathcal{H}[\psi; \zeta]$ should not depend on the electromagnetic gauge. Suppose that one of the variations of ψ^I in the arguments of the symplectic current density $\omega_{cde}(\psi; \delta_1 \psi, \delta_2 \psi)$, say $\delta_2 \psi^I$, is a pure gauge transformation as $\delta_2 g_{ab} = 0$ and $\delta_2 A_a = \nabla_a \chi$, and the other is given by the Lie derivatives along an asymptotic Killing vector ζ^a as $\delta_1 g_{ab} = \mathcal{L}_\zeta g_{ab}$ and $\delta_1 A_a = \mathcal{L}_\zeta A_a$. Then, the variation $\delta_2 \mathcal{H}[\psi; \zeta]$ of $\mathcal{H}[\psi; \zeta]$ under the gauge transformation δ_2 is calculated, by using (4.1) and the Maxwell equation, as

$$\delta_2 \mathcal{H}[\psi; \zeta] = \frac{1}{8\pi} \int_{\partial\mathcal{C}} \epsilon_{bacd} \chi \left[\mathcal{L}_\zeta F^{ba} + \frac{1}{2} g^{de} (\mathcal{L}_\zeta g_{de}) F^{ba} \right]. \quad (4.17)$$

Although $\mathcal{L}_\zeta g_{ab} = \delta_\zeta g_{ab}$ vanishes on the horizon sphere $\partial\mathcal{C}$, due to the boundary condition (4.13), $\mathcal{L}_\zeta F^{ab}$ does not, in general. If we impose $\chi = O(u)$, however, we have $\delta_2 \mathcal{H}[\psi; \zeta] = 0$, as required. Thus, we find, also from gauge invariance of the conserved charges, that the boundary condition (4.15) is reasonable.

While the boundary condition (4.15) on A_a has been determined by incorporating the electromagnetic field into the asymptotic symmetries and imposing the integrability condition of the conserved charges, it does not necessarily ensure that this boundary condition is preserved under the actions of the asymptotic Killing vectors. When we take the Lie derivatives of A_a along the asymptotic Killing vectors ζ^a , we indeed find

$$\delta_\zeta A_a = \mathcal{L}_\zeta A_a = \xi_{(h)}^b A_b \nabla_a \omega + \nabla_a \left[\omega \xi_{(h)}^b \nabla_b (u \chi') + a^{(i)} \xi_{(i)}^b \nabla_b (u \chi') - \lambda A_b \nabla^b \omega \right] + O(u), \quad (4.18)$$

from (3.30) and (4.16). We see that the second term in (4.18) is the gradient of a scalar of $O(u)$, because we have $\xi_{(h)}^b \nabla_b u = O(u)$ and $\xi_{(i)}^b \nabla_b u = O(u)$. Then, by noting that (4.15) yields $n^a \delta A_a = O(u)$ for an arbitrary vector n^a tangent to $u = 0$, a necessary condition for $\delta_\zeta A_a$ to take the form of (4.15) is found as

$$\xi_{(h)}^b A_b n^a \nabla_a \omega = O(u). \quad (4.19)$$

In order that any configurations of A_a achieved by the asymptotic symmetry transformations reside in the covariant phase space, we thus impose either

$$n^a \nabla_a \omega = O(u), \quad (4.20)$$

or

$$\xi_{(h)}^b A_b = O(u). \quad (4.21)$$

When (4.21) is satisfied, (4.18) is consistent with the boundary condition (4.15) for arbitrary ω , and hence the actions of all the asymptotic Killing vectors are appropriately represented in the covariant phase space in this case. If (4.20) is satisfied, however, ω must be constant on $u = 0$. Moreover, as we have seen from (3.29), dependence of ω on u cancels out up to $O(u^2)$, which occurs also in (4.18). Thus, ω in this case must be given as $\omega = \text{const.} + O(u^2)$, and then the asymptotic symmetry group is nothing more than the isometry group. On the other hand, from (4.18), we have $\delta_\zeta A_a = O(u)$ when $\omega = \text{const.} + O(u^2)$, which indicates that the isometry group is represented appropriately in the covariant phase space under the boundary condition (4.15). Therefore, we see that the asymptotic symmetry group is necessarily reduced to the isometry group in the covariant phase space, when the condition (4.21) is not satisfied and hence (4.20) must be imposed.

4.3. Degeneracy

In the case of the non-extreme Kerr–Newman black hole, the regular gauge potential $\bar{A}_a^N$ satisfies (2.16), and hence (4.21) follows from (4.16). Indeed, whenever the surface gravity is non-vanishing, (4.21) is shown to be satisfied on an arbitrary Killing horizon in a regular gauge that satisfies the boundary condition (4.15). A Killing horizon with the non-vanishing surface gravity has a bifurcation surface [22], where the Killing vector $\xi_{(h)}^a$ vanishes, and so does $\xi_{(h)}^b A_b$. In addition, since the boundary condition (4.15) implies

$$\mathcal{L}_{\xi_{(h)}} A_a = \delta_{\xi_{(h)}} A_a = \nabla_a(u \chi') + O(u), \quad (4.22)$$

we have

$$\mathcal{L}_{\xi_{(h)}} \xi_{(h)}^a A_a = O(u). \quad (4.23)$$

Therefore, $\xi_{(h)}^a A_a$ vanishes all over the horizon $u = 0$, and (4.21) holds generally on a Killing horizon with the non-vanishing surface gravity. The actions of all the asymptotic Killing vectors are thus represented in the covariant phase space, without reduction of the asymptotic symmetry group.

However, an extreme black hole does not possess a bifurcation surface, and hence the above argument in the case of non-vanishing surface gravity does not apply to an extreme black hole. It is possible even in the extreme case to choose a gauge which imitates that of a non-extreme black hole, in the sense that (4.21) is satisfied, such as $\bar{A}_a^I$. However, there are no reasons to exclude the gauges that do not satisfy (4.21), such as $\bar{A}_a^{II}$, because the Killing vector $\xi_{(h)}^a$ does not vanish anywhere on the horizon, as we see from (2.29). In the latter case, we need to impose (4.20), and hence the asymptotic symmetry group is necessarily reduced to the isometry group. We emphasize again that the gauges that do not satisfy (4.21) reflect faithfully the global geometric structure of the Killing horizon of an extreme black hole, i.e., absence of a bifurcation surface. In other words, the asymptotic symmetry group must be reduced to the isometry group when we respect this global geometric structure particular to an extreme black hole.

Here we recall that an asymptotic Killing horizon [6] was considered as a local geometric structure generalized from the notion of a Killing horizon, and is defined by the pair (H, ξ^a) of a null hypersurface H and its generator ξ^a . The generator ξ^a of an asymptotic Killing horizon is then given by the asymptotic Killing vectors that become null on H , i.e., those with $a^{(i)} = 0$ in (3.30). Since the function ω in (3.30) is arbitrary, there exist infinitely many asymptotic Killing horizons on the common null hypersurface H . In particular, an arbitrary Killing horizon is accompanied by infinitely many asymptotic Killing horizons, and hence this provides degeneracy associated with the local structure of a Killing horizon. However, when the asymptotic symmetry group is reduced to the isometry group, which occurs in the case of an extreme black hole, the degeneracy of the asymptotic Killing horizons disappears, leaving the extreme black hole isolated.

It was also argued in [6], based on the behavior of the acceleration associated with asymptotic Killing horizons, that the Killing horizon of an extreme black hole shows two contrastive aspects. On one hand, the

Killing horizon of an extreme black hole is considered as isolated from the asymptotic Killing horizons on itself, in contrast with the Killing horizon of a non-extreme black hole, which is continuously deformed into asymptotic Killing horizons. On the other hand, infinitely many asymptotic Killing horizons reside on the Killing horizon both of a non-extreme black hole and of an extreme black hole, and hence an extreme black hole carries as many asymptotic Killing horizons as a non-extreme (near-extreme) black hole. The former aspect is manifest when the global structure near the horizon is probed, while the latter is relevant to the local structure. Therefore, an extreme black hole looks isolated when the global structure of its Killing horizon is respected, but it behaves similarly to a non-extreme black hole as far as its local structure is concerned. We also recall that the Euclidean approach to black hole thermodynamics shows that the entropy of an extreme black hole vanishes [10, 11], which implies that there exists only a single microscopic state, while it is shown in string theory to obey the Bekenstein–Hawking formula [7, 8, 9], indicating that an extreme black hole is accompanied by microscopic states as many as a non-extreme black hole. We now see that the behavior of the asymptotic Killing horizons [6], which was argued based on the acceleration and found consistent with both of these reliable results in string theory and the Euclidean approach, is realized also in the covariant phase space. The covariant phase space of an extreme black hole constructed under the boundary condition that imitates a non-extreme black hole has the size comparable with that of a non-extreme black hole, but the boundary condition that reflects faithfully the global geometric structure of an extreme black hole yields the covariant phase space where degeneracy of asymptotic Killing horizons is not allowed.

We now turn to the issue of the algebra defined by the Poisson brackets (4.7). In particular, it is meaningful to see whether a non-vanishing central charge arises in the Poisson brackets algebra, i.e., whether the sub-group of diffeomorphisms described by the asymptotic Killing vectors is represented with an anomaly. Since we have imposed the boundary condition so that the conserved charges $\mathcal{H}[\psi, \zeta]$ conjugate to the asymptotic Killing vectors ζ^a are integrable, we can compute the central term $K[\zeta_1, \zeta_2]$ by (4.8) and (4.9), where $Q^{ba}(\psi; \zeta)$ is found from the Lagrangian (4.10) as

$$Q^{ba}(\psi; \zeta) = \frac{1}{16\pi} \left[\nabla^a \zeta^b - \nabla^b \zeta^a + 4F^{ab} A_d \zeta^d \right]. \quad (4.24)$$

We then substitute (3.30) and (4.24) into (4.9) and use the field equations. We also note the facts that $\xi_{(h)}^a$ and $\xi_{(t)}^a$ are Killing vectors, which satisfy (2.6) and commute with each other, that the horizon sphere $\partial\mathcal{C}$ does not have its own boundaries, and that $\xi_{(h)}^a$ is hypersurface orthogonal. Then, under the boundary condition imposed above, whether the asymptotic symmetry group is reduced ($\omega = \text{const.} + \mathcal{O}(u^2)$) or not ($\xi_{(h)}^a A_a = \mathcal{O}(u)$), a lengthy calculation yields

$$\delta_{\zeta_2} \mathcal{H}[\bar{\psi}; \zeta_1] = 0. \quad (4.25)$$

Therefore, the Poisson brackets algebra of the conserved charges $\mathcal{H}[\psi, \zeta]$ is given by

$$\left\{ \mathcal{H}[\psi; \zeta_1], \mathcal{H}[\psi; \zeta_2] \right\} = \mathcal{H}[\psi; \mathcal{L}_{\zeta_1} \zeta_2] - \mathcal{H}[\bar{\psi}; \mathcal{L}_{\zeta_1} \zeta_2]. \quad (4.26)$$

When we redefine $\mathcal{H}[\psi; \zeta]$ by an additive constant as

$$\mathcal{H}[\psi; \zeta] \rightarrow \mathcal{H}'[\psi; \zeta] \equiv \mathcal{H}[\psi; \zeta] - \mathcal{H}[\bar{\psi}; \zeta], \quad (4.27)$$

which is achieved by adjusting the integration constant $\mathcal{H}_0[\zeta]$ in (4.5), we see that (4.26) is rewritten as

$$\left\{ \mathcal{H}'[\psi; \zeta_1], \mathcal{H}'[\psi; \zeta_2] \right\} = \mathcal{H}'[\psi; \mathcal{L}_{\zeta_1} \zeta_2]. \quad (4.28)$$

This shows that the central charge vanishes, as in the case of [14], and hence that there arises no anomaly of diffeomorphism invariance. In particular, while the Lie brackets algebra of the asymptotic Killing vectors ζ^a contains the $\text{diff}(S^1)$ or $\text{diff}(R^1)$ sub-algebra [6], the corresponding Poisson brackets sub-algebra does not possess a non-vanishing central charge.

We note that the vanishing central charge in the Poisson brackets algebra is an immediate consequence of (4.25). However, we can argue that (4.25) serves also as an evidence that the microscopic states of black hole thermodynamics are described by the asymptotic Killing horizons. To see this, we set as

$$\zeta_1^a = \xi_{(h)}^a, \quad \zeta_2^a = \xi^a \equiv \omega \xi_{(h)}^a - \lambda \nabla^a \omega + O(u), \quad (4.29)$$

and substitute into (4.25). We then have

$$\delta_\xi \mathcal{H}[\bar{\psi}; \xi_{(h)}] = 0. \quad (4.30)$$

Since the variation δ is defined to act only on the dynamical field variables ψ^I , as we see from the variation of the Lagrangian (4.3), but not on $\xi_{(h)}^a$, (4.30) indicates that the conserved charge $\mathcal{H}[\bar{\psi}; \xi_{(h)}]$ remains unaffected under the infinitesimal transformations

$$\psi^I \rightarrow \psi^I + \mathcal{L}_\xi \psi^I, \quad \xi_{(h)}^a \rightarrow \xi_{(h)}^a, \quad (4.31)$$

i.e., the transformations where the dynamical field variables ψ^I are transformed by the Lie derivatives but the vector $\xi_{(h)}^a$ is fixed. However, we see that the transformations (4.31) are equivalent, up to diffeomorphisms, to the transformations where ψ^I are fixed but $\xi_{(h)}^a$ is transformed as

$$\psi^I \rightarrow \psi^I, \quad \xi_{(h)}^a \rightarrow \xi_{(h)}^a - \mathcal{L}_\xi \xi_{(h)}^a, \quad (4.32)$$

because the transformations (4.31) followed by the infinitesimal diffeomorphisms

$$\psi^I \rightarrow \psi^I - \mathcal{L}_\xi \psi^I, \quad \xi_{(h)}^a \rightarrow \xi_{(h)}^a - \mathcal{L}_\xi \xi_{(h)}^a, \quad (4.33)$$

reduce to (4.32). On the other hand, $\mathcal{H}[\bar{\psi}; \xi_{(h)}]$ (with $\mathcal{H}_0[\xi_{(h)}] = 0$) has been shown [20] to coincide with TS on a stationary and axisymmetric black hole spacetime, where T and S denote the temperature and the entropy of the black hole, respectively. The entropy of the Kerr–Newman black hole is given by the Bekenstein–Hawking formula, and the Lie derivative of the volume element of a horizon sphere vanishes along the null direction on the horizon. Since the zeroth law holds and ξ^a is null on the horizon, we then find that $\mathcal{H}[\bar{\psi}; \xi_{(h)}] = TS$ is invariant under the diffeomorphisms along ξ^a , and hence that it is left unchanged also under the transformations (4.32). Furthermore, we can write [6] as

$$\xi_{(h)}^a - \mathcal{L}_\xi \xi_{(h)}^a = \omega' \xi_{(h)}^a - \lambda \nabla^a \omega' + O(u), \quad (4.34)$$

where $\omega' = 1 + \xi_{(h)}^a \nabla_a \omega$, and thus $\xi_{(h)}^a - \mathcal{L}_\xi \xi_{(h)}^a$ is found to be an asymptotic Killing vector that becomes null on the horizon. As we mentioned above, such an asymptotic Killing vector generates an asymptotic Killing horizon on the same null hypersurface as the Killing horizon. Therefore, the Killing horizon is transformed into asymptotic Killing horizons under the transformations (4.32), and thus we see that the value of the conserved charge $\mathcal{H}[\bar{\psi}; \xi_{(h)}]$ is left unchanged under these transformations from the Killing horizon into the asymptotic Killing horizons. Since $\mathcal{H}[\bar{\psi}; \xi_{(h)}]$ gives the thermodynamical quantity and the function ω in (4.29) is arbitrary, it then implies that all the asymptotic Killing horizons infinitesimally transformed from the Killing horizon are associated with the same macroscopic (thermodynamical) state. This may suggest that the asymptotic Killing horizons are regarded as degenerate from a macroscopic point of view, while they are physically distinguishable [6], as we mentioned in Introduction.

5. Summary and discussion

We first derived the general form of the asymptotic Killing vectors on the Killing horizon of the Kerr–Newman black hole, which was actually shown to be possessed universally by arbitrary Killing horizons [6]. One might suspect that the asymptotic symmetries generated by these asymptotic Killing vectors are nothing more than a sort of gauge, but it should be emphasized that asymptotic Killing horizons, which are described by a sub-group of these asymptotic symmetries, are distinguished physically by the acceleration associated with them [6]. It is then natural to expect that asymptotic Killing horizons have something to do with the universal physics of a Killing horizon, in particular, its thermal feature.

We then considered the covariant phase space associated with these asymptotic Killing vectors. By incorporating the electromagnetic field into the asymptotic symmetries, we found that two types of the boundary condition on the gauge potential are possible in the case of an extreme black hole, and that they give the covariant phase spaces of an extreme black hole with different sizes. The boundary condition that imitates a non-extreme black hole provides the covariant phase space of the size comparable with a non-extreme black hole, while the covariant phase space resulting from the boundary condition that respects the global structure of a Killing horizon particular to an extreme black hole is small enough to exclude the degeneracy of asymptotic Killing horizons. It should be stressed, however, that what plays an essential role is not the electromagnetic gauge, but the geometric structure of the Killing horizon of an extreme black hole. The electromagnetic field simply probes this geometric structure. Even without the electromagnetic field, as in the case of the Kerr black hole, it is actually possible to reduce the covariant phase space of the asymptotic Killing vectors by imposing stringent conditions that respect the global geometric structure of a Killing horizon particular to an extreme black hole. Therefore, the behavior of the asymptotic Killing horizons of an extreme black hole, which was found from the acceleration associated with them [6], is properly realized also in the covariant phase space. In particular, the covariant phase space of the asymptotic Killing vectors is consistent with both the entropy of an extreme black hole in string theory [7, 8, 9] and that in the Euclidean approach [10, 11], which are apparently conflicting with each other but might be viewed as dual aspects of a single phenomenon.

It is important also to mention the issue of the central charge in the Poisson brackets algebra, which represents an anomaly of diffeomorphism invariance. We showed in this paper that the central charge vanishes. Thus, straightforward application of the same method as that applied to the *infinity* of the B.T.Z. black hole spacetime [3] does not reproduce the Bekenstein–Hawking formula. One might then consider that this result indicates that the asymptotic symmetries analyzed in this paper have nothing to do with microscopic states responsible for the thermal feature of a horizon, particularly if we persist in following the derivation in the case of the B.T.Z. black hole [3]. However, there actually exists an explicit *microscopic* model in the theory of induced gravity [24], where the Bekenstein–Hawking formula has been reproduced based on the asymptotic symmetries with the *vanishing total central charge*, while each microscopic field in this model carries a non-vanishing central charge. Moreover, in the recent approach to Hawking radiation [25], it was found [26] that the anomaly cancellation condition, which thus requires that *anomalies of diffeomorphism and gauge invariance are absent*, along with appropriate conditions on the effective energy-momentum, lead to the Hawking flux with the correct temperature. Therefore, the vanishing central charge looks consistent with the thermal feature of a horizon, at least in an effective theory. Furthermore, we saw that the vanishing central charge provides an evidence that all the asymptotic Killing horizons on a Killing horizon are associated with the same macroscopic state, which may imply that the microscopic states of black hole thermodynamics are described by the asymptotic Killing horizons. Probably, not all of these asymptotic Killing horizon will contribute to black hole entropy, and a sort of quantization condition may be imposed in a quantum theory, which will allow only a discrete subset of the asymptotic Killing horizons to constitute a thermal object. It will be interesting to analyze this issue further in future investigations.

Acknowledgments

I would like to thank A. Hosoya, K. Maeda, M. Natsuume, G. Kang and M. Park for useful discussions and encouragement.

References

- [1] Bardeen J M, Carter B and Hawking S W 1973 Commun. Math. Phys. **31** 161
- [2] Hawking S W 1975 Commun. Math. Phys. **43** 199
- [3] Strominger A 1998 J. High Energy Phys. **02** 009

- [4] Fursaev D V 2005 *Phys. Part. Nucl.* **36** 81 (*Preprint* gr-qc/0404038)
- [5] Carlip S 2006 *Preprint* gr-qc/0601041
- [6] Koga J 2006 *Preprint* gr-qc/0604054
- [7] Strominger A and Vafa C 1996 *Phys. Lett. B* **379** 99
- [8] Horowitz G T 1996 *Preprint* gr-qc/9604051
- [9] Sen A 2005 *JHEP* **09** 038
- [10] Hawking S W, Horowitz G T and Ross S F 1995 *Phys. Rev. D* **51** 4302
- [11] Teitelboim C 1995 *Phys. Rev. D* **51** 4315
- [12] Bondi H, van der Burg M G J and Metzner A W K 1962 *Proc. Roy. Soc. (London)* **A269** 21; Sachs R K 1962 *Proc. Roy. Soc. (London)* **A270** 103; Sachs R K 1962 *Phys. Rev.* **128** 2851
- [13] Brown J D and Henneaux M 1986 *Comm. Math. Phys.* **104** 207
- [14] Koga J 2001 *Phys. Rev. D* **64** 124012
- [15] Silva S 2002 *Class. Quantum Grav.* **19** 3947
- [16] Barnich G and Brandt F 2002 *Nucl. Phys. B* **633** 3
- [17] Frolov V P and Novikov I D 1998 *Black Hole Physics* (The Netherlands: Kluwer Academic Publishers)
- [18] Wald R M 1994 *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics* (Chicago: The University of Chicago Press)
- [19] Lee J and Wald R M 1990 *J. Math. Phys.* **31** 725
- [20] Wald R M 1993 *Phys. Rev. D* **48** 3427; Iyer V and Wald R M 1994 *Phys. Rev. D* **50** 846
- [21] Wald R M and Zoupas A 2000 *Phys. Rev. D* **61** 084027
- [22] Ratz I and Wald R M 1992 *Class. Quantum Grav.* **9** 2643; Ratz I and Wald R M 1996 *Class. Quantum Grav.* **13** 539
- [23] Henneaux M and Teitelboim C 1985 *Commun. Math. Phys.* **98** 391
- [24] Frolov V, Fursaev D and Zelnikov A 2003 *JHEP* **03** 038
- [25] Robinson S P and Wilczek F 2005 *Phys. Rev. Lett.* **95** 011303
- [26] Iso S, Umetsu H and Wilczek F 2006 *Phys. Rev. Lett.* **96** 151302; Iso S, Umetsu H and Wilczek F 2006 *Phys. Rev. D* **74** 044017